

Temperature Effects on Efficacy of *Choetospila elegans* (Hymenoptera: Pteromalidae) to Suppress *Rhyzopertha dominica* (Coleoptera: Bostrichidae) in Stored Wheat

P. W. FLINN

Grain Marketing and Production Research Center, USDA-ARS, Manhattan, KS 66502

J. Econ. Entomol. 91(1): 320-323 (1998)

ABSTRACT Laboratory studies were conducted to assess the effectiveness of the parasitoid wasp *Choetospila elegans* (Westwood) for controlling *Rhyzopertha dominica* (F.), lesser grain borer, in wheat at 32 and 25°C. The 2 temperature regimes were used to simulate an unaerated bin of wheat and a bin aerated at harvest time. Two adult male and 2 adult female *R. dominica* were each released into containers with 19 kg of hard red winter wheat. An equal number of adult *C. elegans* were released into half of the containers. Half the containers were kept at 25°C and half at 32°C. Suppression of *R. dominica* population growth by *C. elegans* was much greater at 25 than at 32°C. After 161 d, *R. dominica* density in the containers with *C. elegans* was 9,185/kg at 32°C, and 10/kg at 25°C. At 25°C, *C. elegans* was able to locate and parasitize most of the larvae that were produced by the adult beetles. This resulted in a very high level of population suppression (99% in comparison to the control at 25°C). In contrast, at 32°C, beetle suppression was only 50% in comparison to containers without *C. elegans* at this temperature. This study suggests that when augmentative parasite releases are made with *C. elegans*, better host suppression would be achieved by cooling the grain to 25°C shortly after harvest, rather than leaving it unaerated for the summer.

KEY WORDS *Rhyzopertha dominica*, *Choetospila elegans*, biological control, stored-products, temperature.

BIOLOGICAL CONTROL IS an overlooked component of integrated pest management of stored grain. Most of the parasitoids that attack the primary beetle pests are in the families Pteromalidae and Bethyliidae (Hagstrum and Flinn 1992). These hymenopterous parasitoids are very small (1-2 mm) and do not feed on the grain. They will normally die or leave the grain within 5-10 d if no beetles are present in the grain. These parasitoids are found naturally in stored grain, which suggests that once released they may continue to suppress pests for many years (Sinha et al. 1979). Because the adult wasps are external to the grain, they can be removed easily from it using normal cleaning processes. The lesser grain borer, *Rhyzopertha dominica* (F.), is one of the more common and damaging insect pests of stored wheat in the United States (Storey et al. 1984). Adults feed primarily on the wheat endosperm and cause considerable damage. *R. dominica* larvae develop within the grain kernel and cannot be removed from the grain by normal cleaning procedures.

Choetospila elegans (Westwood) is a small pteromalid wasp that attacks the coleopterans *R. dominica*, *Sitophilus* spp., *Stegobium paniceum* (L.), *Callosobruchus* spp., and the lepidopteran, *Sitotroga cerealella* (Oliver) (Burks 1979). This wasp normally parasitizes larvae that are feeding inside the

grain kernel. Although wasp larvae can complete development on 3rd instar and prepupal *R. dominica*, larval survivorship is highest when laid on 4th-instar *R. dominica* (P. Flinn, unpublished data). They normally lay 1 egg externally on each host (Sharifi 1972). If >1 egg is present, only 1 larva completes development. At 32°C, it takes ≈15 d to complete development on *R. dominica* (unpublished data). The generation time of this wasp is about half that of *R. dominica*. If hosts are available, female wasps live for 10-20 d at 32°C. A single female *C. elegans* can parasitize up to 6 *R. dominica* per day (unpublished data). In a previous field study, Flinn et al. (1996) showed that *C. elegans* was very effective in suppressing *R. dominica* in 27 ton bins of stored wheat. *R. dominica* populations were suppressed by >91% in relation to control bins.

Biological control can be more effective when used in conjunction with other control methods (Van Driesche and Bellows 1996). In stored grain, one of the more effective nonchemical control methods is to cool the grain with aeration fans. Aeration, using electric powered fans, can be used to cool the grain earlier; thus it suppresses insect population growth sooner in the storage period (Flinn et al. 1997). The Kansas State extension program has recently begun advocating early aeration (starting at harvest) with automatic aeration con-

trollers as the best nonchemical insect suppression method for stored grain. A study conducted in Kansas showed that early aeration, using automatic fan controllers, allowed safe storage of grain for many months (Reed and Harner 1997). A controller turns on the fan when the outside air temperature is low enough to cool the grain efficiently. Compared with manual aeration, automatic aeration controllers cool the grain sooner and with less moisture loss, which is an important consideration with stored wheat. Under typical summer conditions in Kansas, automatic aeration controllers will reduce the grain temperature from 32°C down to ≈25°C (Reed and Harner 1997). The population growth rate of *R. dominica* is about half as fast at 25°C than at 32°C (Hagstrum and Milliken 1988).

Previous biological control studies in stored grain have not investigated the combined effects of parasitoid wasps and cool grain temperatures. In this study, I investigated the effects of 2 temperatures, which simulated unaerated and summer aeration of wheat, on the ability of *C. elegans* to suppress *R. dominica* population growth.

Materials and Methods

Experiments were conducted in 12 cylindrical plastic pails (22.7 liter, 38 cm tall by 31 cm diameter). The containers had tight fitting snap-on lids, with a 21-cm hole covered with 122-mesh polyester silk-screen (Majestic Arts, Somers, NY). The containers were filled with 19 kg of hard red winter wheat (12% moisture). Two adult female and 2 adult male *R. dominica* beetles were placed in each container. The adults were obtained from a laboratory culture and were ≈1 wk old. Six of the containers were placed in an environmental chamber maintained at 32 ± 1°C and 65 ± 10% RH, and 6 were placed in an environmental chamber maintained at 25 ± 1°C and 65 ± 10% RH. After 20 d, 2 adult male and 2 adult female *C. elegans* wasps were placed in 3 of the containers in the 32°C chamber. The wasps were obtained from a laboratory culture and were ≈2 d old. Ten days later, 2 adult male and 2 adult female *C. elegans* were placed in 3 of the containers in the 25°C chamber. *C. elegans* prefers to parasitize 4th-instar *R. dominica*. Therefore, the parasitoids were released 10 d later in the 25°C than in the 32°C chambers so that stadiums 1–4 would be available in containers in both chambers, and so that few pupae would have developed yet.

Sixty-three days from the initial beetle release, 9 grain samples, each ≈45 g, were taken in each container using a miniature grain trier (Flinn and Hagstrum 1995). The containers were sampled every 2 wk up to 161 d from the initial beetle release. The samples were sieved and insects counted.

Differences between means of *R. dominica* density were tested using a t-test ($\alpha = 0.05$) (SAS Institute 1990).




Fig. 1. Mean *R. dominica* density in containers with 19 kg of wheat with and without *C. elegans* added at 25 and 32°C. Vertical bars indicate standard errors of the mean.

Results and Discussion

Suppression of *R. dominica* population growth by *C. elegans* was much greater at 25 than at 32°C. After 161 d of storage, *R. dominica* density in the containers with *C. elegans* was 9,185/kg at 32°C, and 10/kg at 25°C (Fig. 1). At 32°C, there were significant differences in mean *R. dominica* density between containers with and without *C. elegans* at each sampling date. At 25°C, there were significant differences in mean *R. dominica* density between containers with and without *C. elegans* at each sampling date, except for the first 2 sampling dates. Population growth of *R. dominica* was also suppressed by lower temperature alone. After 161 d of storage in containers without *C. elegans*, *R. dominica* density was significantly greater in containers held at 32°C than at 25°C (18,673/kg at 32°C and 1,010/kg at 25°C).

Choetospila elegans reached greater densities in the containers held at 32°C than at 25°C (Fig. 2). At 32°C, *C. elegans* reached a peak of 20 insects per kilogram after 91 d. However, after 133 d, very few *C. elegans* were found in the 32°C containers. After 105 d at 32°C, beetle density in the containers with *C. elegans* reached an average of 616/kg. This is a




Fig. 2. Mean *C. elegans* density in containers with 19 kg of wheat at 25 and 32°C. Vertical bars indicate standard errors of the mean.

very high beetle density, and may have suppressed wasp population growth by interfering with parasitization. At 25°C, *C. elegans* increased steadily to a maximum density of 3/kg after 146 d.

In general, the percentage of reduction in *R. dominica* density caused by *C. elegans* was greater at 25°C than at 32°C (Fig. 3). After 133 d, percentage reduction decreased to ≈50% in the containers held at 32°C and increased to 99% in the containers held at 25°C treatment. At 25°C, *C. elegans* may have been able to locate and parasitize most of the larvae that were produced by the original *R. dominica* adults, and this resulted in a very high level of population suppression. In contrast, at 32°C, *C. elegans* may have been unable to locate and parasitize most of the beetle larvae. At the warmer temperature, the beetles had a higher oviposition and developmental rate, and thus the number of beetle larvae *C. elegans* would need to parasitize would be much higher to prevent *R. dominica* population growth. Because *C.*




Fig. 3. Percentage reduction of *R. dominica* at 25 and 32°C caused by *C. elegans* in relation to the absence of *C. elegans* at each temperature.

elegans does not attack adult beetles, the ability of *C. elegans* to suppress the population would diminish as the number of unparasitized *R. dominica* larvae that developed into adults increased. Although *C. elegans* population growth rate is also higher at 32°C, their ability to find and parasitize most of the 4th-instar *R. dominica* larvae may not be greater at the higher temperature. If we had released more *C. elegans* at the start of the experiment, they may have suppressed the beetle population more at the higher temperature. However, the point of this experiment was to compare wasp suppression ability at the 2 temperatures and not over a range of wasp release densities.

In the United States, fall aeration is often started in October. Wheat that is not aerated at harvest (early July in Kansas) may not be aerated for 80–100 d, and it may take until November until the temperature of the grain is below that at which *R. dominica* can develop (<20°C) (Fields and Muir 1996). In the containers held at 32°C with *C. elegans*, there were 331 *R. dominica* per kilogram after 91 d. At 25°C, there were 3/kg in the containers with *C. elegans*, and 11 *R. dominica* per kilogram in the containers without *C. elegans* after 91 d. In the southern United States, where it is difficult to cool grain below 25°C, the expected *R. dominica* densities for the respective treatments may be similar to the densities at day 161 of this experiment.

When augmentative parasite releases are made with *C. elegans*, this study suggests that better *R. dominica* suppression would be achieved by cooling the grain with aeration as soon as it is put in the bin. Even when augmentative releases are not used, reducing the grain temperature with early aeration will suppress *R. dominica* because of decreased population growth at lower temperatures. In addition, because natural populations of *C. elegans* are often found in grain bins, lower temperatures will help these natural populations to suppress *R. dominica*.

Acknowledgments

I thank T. Phillips (Department of Entomology, Oklahoma State University), B. Dover (Department of Entomology, Kansas State University), and R. Howard (Grain Marketing and Production Research Center, USDA-ARS) for reviewing an earlier version of the manuscript, and Ken Friesen for excellent technical support.

References Cited

- Burks, D. B. 1979. Family Pteromalidae, pp. 768–834. In V. Krombein, P. D. Hurd, D. R. Smith, and D. B. Burks [eds.], *Catalog of Hymenoptera in America north of Mexico*. Smithsonian Institution, Washington, DC.
- Fields, P. G., and W. E. Muir. 1996. Physical control, pp. 195–221. In B. Subramanyam and D. W. Hagstrum [eds.], *Integrated management of insects in stored products*. Marcel Dekker, New York.
- Flinn, P. W., and D. W. Hagstrum. 1995. Simulation model of *Cephalonomia waterstoni* (Hymenoptera: Bethyloidea) parasitizing the rusty grain beetle (*C.*

- leoptera: Cucujidae). Environ. Entomol. 24: 1608–1615.
- Flinn, P. W., D. W. Hagstrum, and W. H. McGaughey.** 1996. Suppression of beetles in stored wheat by augmentative releases of parasitic wasps. Environ. Entomol. 25: 505–511.
- Flinn, P. W., D. W. Hagstrum, and W. E. Muir.** 1997. Effects of time of aeration, bin size, and latitude on insect populations in stored wheat: a simulation study. J. Econ. Entomol. 90: 646–651.
- Hagstrum, D. W., and P. W. Flinn.** 1992. Integrated pest management of stored-grain insects, pp. 535–562. In D. B. Sauer [ed.], Storage of cereal grains and their products. American Association of Cereal Chemists, St. Paul, MN.
- Hagstrum, D. W., and G. A. Milliken.** 1988. Quantitative analysis of temperature, moisture, and diet factors affecting insect development. Ann. Entomol. Soc. Am. 81: 539–546.
- Reed, C., and J. Harner.** 1997. Thermostatically controlled aeration for insect control in stored hard red winter wheat. J. Appl. Engineering. (in press).
- SAS Institute.** 1990. SAS/STAT user's guide, version 6, 4th ed. SAS Institute, Cary, NC.
- Sharifi, S.** 1972. Radiographic studies of the parasite *Choetospila elegans* on the maize weevil, *Sitophilus zeamais*. Ann. Entomol. Soc. of Am. 65: 852–856.
- Sinha, R. N., H.A.H. Wallace, B. Reiser, and L. P. Lefkovich.** 1979. Interrelations of arthropods and microorganisms in damp bulk stored wheat—a multivariate study. Res. Popul. Ecol. 21: 40–67.
- Storey, C. L., D. B. Sauer, and D. Walker.** 1984. Present use of pest management practices in wheat, corn, and oats stored on the farm. J. Econ. Entomol. 77: 784–788.
- Van Driesche, R. G. and T. S. Bellows.** 1996. Biological control. Chapman & Hall, New York.

Received for publication 6 June 1997; accepted 27 October 1997.
